arXiv:1702.01960v1 [math.CA] 7 Feb 2017

CERTAIN NEW UNIFIED INTEGRALS ASSOCIATED WITH THE PRODUCT OF GENERALIZED STRUVE FUNCTION

KOTTAKKARAN SOOPPY NISAR

Abstract. We aim to present two new generalized integral formulae involving product of generalized Struve function $\mathcal{W}_{p,b,c}(z)$, which are expressed in terms of the generalized Lauricella functions. The main results presented here, being very general character, reduce to yield known and new integral formulae. Some special cases of our main results are also considered.

1. Introduction and Preliminaries

A solution of the following non-homogeneous Bessel's differential equation

$$x^2y''(x)+xy'(x)+\left(x^2-p^2\right)y(x)=\frac{4\left(x/2\right)^{\alpha+1}}{\sqrt{\pi}\,\Gamma\left(\alpha+1/2\right)}\quad(\alpha\in\mathbb{C}) \tag{1.1}$$

is known as the Struve function $\mathcal{H}_\alpha$ of order α, where Γ is the familiar gamma function (see, e.g., [18, Section 1.1]). Here and in the following, let $\mathbb{C}$, $\mathbb{R}^+$, and $\mathbb{N}$ be the sets of complex numbers, positive real numbers, and positive integers, respectively, and let $\mathbb{N}_0:=\mathbb{N}\cup\{0\}$. The Struve functions occur in many areas such as water wave and surface-wave problems (see [2, 9]), problems on unsteady aerodynamics [16], particle quantum dynamical studies of spin decoherence [15] and nanotubes [14]. The Struve functions $\mathcal{H}_v(z)$ and $\mathcal{L}_v(z)$ are defined as the following infinite series

$$\mathcal{H}_v(z)=\left(\frac{z}{2}\right)^{v+1}\sum_{k=0}^{\infty}\frac{(-1)^k}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+v+\frac{1}{2}\right)}\left(\frac{z}{2}\right)^{2k} \tag{1.2}$$

and

$$\mathcal{L}_v(z)=\left(\frac{z}{2}\right)^{v+1}\sum_{k=0}^{\infty}\frac{1}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+v+\frac{1}{2}\right)}\left(\frac{z}{2}\right)^{2k}. \tag{1.3}$$

The generalized Struve function is given as follows (see, e.g., [11]):

$$\mathcal{W}_{p,b,c}(z)=\sum_{k=0}^{\infty}\frac{(-c)^k\left(\frac{z}{2}\right)^{2k+p+1}}{\Gamma\left(k+\frac{3}{2}\right)\Gamma\left(k+p+\frac{b+2}{2}\right)}\quad(p,\,b,\,c\in\mathbb{C}). \tag{1.4}$$

2000 *Mathematics Subject Classification.* Primary 33B20, 33C20; Secondary 33B15 ,33C05.

Key words and phrases. Pochhammer symbol, Gamma function, Generalized hypergeometric function ${}_pF_q$, Generalized (Wright) hypergeometric functions ${}_p\Psi_q$, generalized Struve function and Oberhettinger integral formula.

An interesting further generalization of the generalized hypergeometric series ${}_pF_q$ (see, e.g., [18, Section 1.5]) is due to Fox [7] and Wright [21, 22] who studied the asymptotic expansion of the generalized (Wright) hypergeometric function defined by (see [20, p. 21])

$$ {}_p\Psi_q\left[\begin{matrix}(\alpha_1,\,A_1),\,\ldots,\,(\alpha_p,\,A_p);\\(\beta_1,\,B_1),\,\ldots,\,(\beta_q,\,B_q);\end{matrix}\,z\right]=\sum_{k=0}^{\infty}\frac{\prod\limits_{j=1}^{p}\Gamma\left(\alpha_j+A_j\,k\right)}{\prod\limits_{j=1}^{q}\Gamma\left(\beta_j+B_j\,k\right)}\,\frac{z^k}{k!}, \tag{1.5}$$

where the coefficients $A_1,\,\ldots,\,A_p\in\mathbb{R}^+$ and $B_1,\,\ldots,\,B_q\in\mathbb{R}^+$ such that

$$1+\sum_{j=1}^{q}B_j-\sum_{j=1}^{p}A_j\geqq 0. \tag{1.6}$$

A special case of (1.5) is

$$ {}_p\Psi_q\left[\begin{matrix}(\alpha_1,\,1),\,\ldots,\,(\alpha_p,\,1);\\(\beta_1,\,1),\,\ldots,\,(\beta_q,\,1);\end{matrix}\,z\right]=\frac{\prod\limits_{j=1}^{p}\Gamma\left(\alpha_j\right)}{\prod\limits_{j=1}^{q}\Gamma\left(\beta_j\right)}\,{}_pF_q\left[\begin{matrix}\alpha_1,\,\ldots,\,\alpha_p;\\\beta_1,\,\ldots,\,\beta_q;\end{matrix}\,z\right]. \tag{1.7}$$

An interesting further several-variable-generalization of the generalized Lauricella series (see, for example, [20, p. 36, Eq. (19)]) is defined by (*cf.* Srivastava and Daoust [19, p. 454]; see also [20, p. 37])

$$\begin{aligned}F_{C:D^{(1)};\cdots;D^{(n)}}^{A:B^{(1)};\cdots;B^{(n)}}\left(\begin{array}{c}z_1\\\vdots\\z_n\end{array}\right)&=F_{C:D^{(1)};\cdots;D^{(n)}}^{A:B^{(1)};\cdots;B^{(n)}}\left(\begin{matrix}[(a):\theta^{(1)},\ldots,\theta^{(n)}]:\\[(c):\psi^{(1)},\ldots,\psi^{(n)}]:\end{matrix}\right.\\&\qquad\left.\begin{matrix}[(b)^{(1)}:\phi^{(1)}];\ldots;[(b)^{(n)}:\phi^{(n)}];\\[(d)^{(1)}:\delta^{(1)}];\ldots;[(d)^{(n)}:\delta^{(n)}];\end{matrix}\,z_1,\ldots,z_n\right)\\&=\sum_{k_1,\ldots,k_n=0}^{\infty}\Omega(k_1,\ldots,k_n)\,\frac{z_1^{k_1}}{k_1!}\cdots\frac{z_n^{k_n}}{k_n!},\end{aligned} \tag{1.8}$$

where, for convenience,

$$\Omega(k_1,\ldots,k_n)=\frac{\prod\limits_{j=1}^{A}(a_j)_{k_1\theta_j^{(1)}+\cdots+k_n\theta_j^{(n)}}\prod\limits_{j=1}^{B^{(1)}}(b_j^{(1)})_{k_1\phi_j^{(1)}}\cdots\prod\limits_{j=1}^{B^{(n)}}(b_j^{(n)})_{k_n\phi_j^{(n)}}}{\prod\limits_{j=1}^{C}(c_j)_{k_1\psi_j^{(1)}+\cdots+k_n\psi_j^{(n)}}\prod\limits_{j=1}^{D^{(1)}}(d_j^{(1)})_{k_1\delta_j^{(1)}}\cdots\prod\limits_{j=1}^{D^{(n)}}(d_j^{(n)})_{k_n\delta_j^{(n)}}}, \tag{1.9}$$

the coefficients

$$\begin{cases} \theta_j^{(m)} \ (j=1,\ldots,A);\ \phi_j^{(m)} \ (j=1,\ldots,B^{(m)}); \\ \psi_j^{(m)} \ (j=1,\ldots,C);\ \delta_j^{(m)} \ (j=1,\cdots,D^{(m)});\ \forall\, m\in\{1,\ldots,n\} \end{cases} \tag{1.10}$$

are real and positive, and (a) abbreviates the array of A parameters $a_1,\ldots,a_A$, $(b^{(m)})$ abbreviates the array of $B^{(m)}$ parameters

$$b_j^{(m)} \quad (j=1,\ldots,B^{(m)}); \quad \forall\, m\in\{1,\ldots,n\},$$

with similar interpretations for (c) and $(d^{(m)})$ $(m=1,\ldots,n)$; *et cetera*.

For more details about the generalized Lauricella function (1.8), the reader may be referred, for example, to [6, 8, 17, 19, 20].

We also need the following integral formula due to Oberhettinger [13]

$$\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}dx = 2\lambda a^{-\lambda}\left(\frac{a}{2}\right)^{\mu}\frac{\Gamma\left(2\mu\right)\Gamma\left(\lambda-\mu\right)}{\Gamma\left(1+\lambda+\mu\right)} \tag{1.11}$$

$$\left(a\in\mathbb{R}^+;\ 0<\Re(\mu)<\Re(\lambda)\right).$$

Many integral formulas involving Bessel functions, generalized Bessel functions, and modified Bessel functions, and so on, have recently been established (see, e.g., [1, 3–5, 10, 12]). In this sequel, we aim to present two new integral formulas involving a finite product of the generalized Struve functions (1.4), which are expressed in terms of the generalized Lauricella function (1.8). The main formulas presented here, being very general, reduce to yield known and new integral formulas. Some interesting special cases are also considered.

2. Main results

Here, we establish two integral formulas involving a finite product of the generalized Struve functions (1.4), which are asserted by Theorems 1 and 2.

Theorem 1. *Let $n\in\mathbb{N}$ be fixed and a, $y_j\in\mathbb{R}^+$ $(j=1,\ldots,n)$. Also, let b, c, p_j, λ, $\mu\in\mathbb{C}$ $(j=1,\ldots,n)$ with $0<\Re(\mu)<\Re\left(\lambda+\mathbf{p}\right)+n$, where*

$$\mathbf{p}:=\sum_{j=1}^{n}p_j. \tag{2.1}$$

Then

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\prod_{j=1}^{n}\mathcal{W}_{p_j,b,c}\left(\frac{y_j}{x+a+\sqrt{x^2+2ax}}\right)dx\\
&=\mathcal{A}(a,\mathbf{p},\lambda,\mu,p_j,y_j:j=1,\ldots,n)\\
&\quad\times F_{2:2;\cdots;2}^{2:1;\cdots;1}\left(\begin{matrix}[1+\lambda+\mathbf{p}+n:2,\ldots,2],[\lambda+\mathbf{p}+n-\mu:2,\ldots,2]:\\ [\lambda+\mathbf{p}+n:2,\ldots,2],[1+\lambda+\mathbf{p}+n+\mu:2,\ldots,2]:\end{matrix}\right.\\
&\qquad\qquad\qquad

we obtain

$$\begin{aligned}\mathcal{L} =&\Lambda(a,\mathbf{p},\lambda,\mu,p_j,y_j : j=1,\dots,n)\\ &\times\sum_{k_1,\dots,k_n=0}^{\infty}\frac{(-c)^{k_1}\,\left(\frac{y_1}{2}\right)^{2k_1}\cdots(-c)^{k_n}\,\left(\frac{y_n}{2}\right)^{2k_n}\,a^{-2(k_1+\cdots+k_n)}}{\left(\frac{3}{2}\right)_{k_1}\left(p_1+\frac{b+2}{2}\right)_{k_1}\cdots\left(\frac{3}{2}\right)_{k_n}\left(p_n+\frac{b+2}{2}\right)_{k_n}}\\ &\times\frac{(1+\lambda+\mathbf{p}+n)_{2(k_1+\cdots+k_n)}\,(\lambda+\mathbf{p}+n-\mu)_{2(k_1+\cdots+k_n)}}{(\lambda+\mathbf{p}+n)_{2(k_1+\cdots+k_n)}\,(1+\lambda+\mathbf{p}+n+\mu)_{2(k_1+\cdots+k_n)}},\end{aligned}$$

which, upon the multiple summations being expressed in terms of (1.8), leads to the right-hand side of (2.2).

□

Theorem 2. *Let $n\in\mathbb{N}$ be fixed and $a,\ y_j\in\mathbb{R}^+$ $(j=1,\dots,n)$. Also, let $b,\ c,\ p_j,\ \lambda,\ \mu\in\mathbb{C}$ $(j=1,\dots,n)$ with $\Re(\mu+\mathbf{p})>-n$ and $\Re(\lambda)>\Re(\mu)$. Then*

$$\begin{aligned}&\int_0^{\infty}x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\prod_{j=1}^{n}\mathcal{W}_{p_j,b,c}\left(\frac{x\,y_j}{x+a+\sqrt{x^2+2ax}}\right)dx\\ &=\mathcal{B}(a,\mathbf{p},\lambda,\mu,p_j,y_j : j=1,\dots,n)\\ &\quad\times F_{2:2;\cdots;2}^{2:1;\cdots;1}\left(\begin{matrix}[2\mu+2\mathbf{p}+2n:4,\dots,4],[1+\lambda+\mathbf{p}+n:2,\dots,2]:\\ [1+\lambda+\mu+2\mathbf{p}+2n:4,\dots,4],[\lambda+\mathbf{p}+n:2,\dots,2]:\end{matrix}\right.\\ &\qquad\left.\begin{matrix}[1:1];\cdots;[1:1];\\ \left[\frac{3}{2}:1\right],\left[p_1+\frac{b+2}{2}:1\right];\cdots;\left[\frac{3}{2}:1\right],\left[p_n+\frac{b+2}{2}:1\right];\end{matrix}\ -\frac{c\,y_1^2}{16},\cdots,-\frac{c\,y_n^2}{16}\right),\end{aligned}\tag{2.6}$$

where $1;\cdots;1$ and $2;\cdots;2$ are n arrays, respectively, and

$$\begin{aligned}&\mathcal{B}(a,\mathbf{p},\lambda,\mu,p_j,y_j : j=1,\dots,n)\\ &:=\frac{(\lambda+\mathbf{p}+n)\,2^{1-\mu-2\mathbf{p}-2n}\,a^{\mu-\lambda}\,\Gamma(\lambda-\mu)\,\Gamma(2\mu+2\mathbf{p}+2n)\prod_{j=1}^{n}y_j^{p_j+1}}{\left\{\Gamma\left(\frac{3}{2}\right)\right\}^n\,\Gamma(1+\lambda+\mu+2\mathbf{p}+2n)\prod_{j=1}^{n}\Gamma\left(p_j+\frac{b+2}{2}\right)},\end{aligned}$$

and $\mathbf{p}$ is the same as in (2.1).

Proof. The proof runs parallel to that of Theorem 1. We omit the details. □

3. Special cases

Here, we consider some special cases of the main results.

Setting $n=1$ in the result in Theorem 1, we have an integral formula involving the generalized Struve function (1.4), which is given in Corollary 3.1.

Corollary 3.1. *Let a, $y \in \mathbb{R}^+$. Also, let b, c, p, λ, $\mu \in \mathbb{C}$ with $0 < \Re(\mu) < \Re(\lambda + p) + 1$. Then*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\mathcal{W}_{p,b,c}\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)dx\\
&=(1+\lambda+p)\,2^{-\mu-p}\,a^{\mu-1-\lambda-p}\,y^{p+1}\,\frac{\Gamma(2\mu)\,\Gamma(1+\lambda+p-\mu)}{\Gamma\left(\frac{3}{2}\right)\,\Gamma(2+\lambda+p+\mu)\,\Gamma\left(1+\frac{b}{2}+p\right)}\\
&\times {}_4F_5\left[\begin{matrix}\frac{3}{2}+\frac{\lambda}{2}+\frac{p}{2},\ \frac{1}{2}+\frac{\lambda}{2}+\frac{p}{2}-\frac{\mu}{2},\ 1+\frac{\lambda}{2}+\frac{p}{2}-\frac{\mu}{2},\ 1;\\ \frac{1}{2}+\frac{\lambda}{2}+\frac{p}{2},\ 1+\frac{\lambda}{2}+\frac{p}{2}+\frac{\mu}{2},\ \frac{3}{2}+\frac{\lambda}{2}+\frac{p}{2}+\frac{\mu}{2},\ 1+\frac{b}{2}+p,\ \frac{3}{2};\end{matrix}\ -\frac{cy^2}{4a^2}\right].
\end{aligned}\tag{3.1}$$

Setting $n = 1$ in the result in Theorem 2, we have another integral formula involving the generalized Struve function (1.4), which is given in Corollary 3.2.

Corollary 3.2. *Let a, $y \in \mathbb{R}^+$. Also, let b, c, p, λ, $\mu \in \mathbb{C}$ with $\Re(\mu + p) > -1$ and $\Re(\lambda) > \Re(\mu)$. Then*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\mathcal{W}_{p,b,c}\left(\frac{xy}{x+a+\sqrt{x^2+2ax}}\right)dx\\
&=2^{-\mu-2p-1}\,a^{\mu-\lambda}\,y^{p+1}\,\Gamma(\lambda-\mu)\\
&\times {}_3\Psi_4\left[\begin{matrix}(1,\,1)\,,\ (\lambda+p+2,\,2)\,,\ (2\mu+2p+2,\,4)\,;\\ \left(\frac{3}{2},\,1\right),\ \left(p+\frac{b+2}{2},\,1\right),\ (\lambda+p+1,\,2)\,,\ (\lambda+\mu+2p+3,\,4)\,;\end{matrix}\ -\frac{c\,y^2}{16}\right].
\end{aligned}\tag{3.2}$$

It is easy to see from (1.2) and (1.4) that

$$\mathcal{W}_{p,-1,1}(z) = \mathcal{H}_p(z). \tag{3.3}$$

Considering the relation (3.3) and setting $n = 1$, $b = -1$, and $c = 1$ in Theorems 1 and 2 (or setting $b = -1$ and $c = 1$ in Corollaries 3.1 and 3.2, we obtain two integral formulas involving the Struve function $\mathcal{H}_p$ of order p, which are given, respectively, in Corollaries 3.3 and 3.4.

Corollary 3.3. *Let a, $y \in \mathbb{R}^+$. Also, let p, λ, $\mu \in \mathbb{C}$ with $0 < \Re(\mu) < \Re(\lambda + p) + 1$. Then*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\mathcal{W}_{p,b,c}\left(\frac{y}{x+a+\sqrt{x^2+2ax}}\right)dx\\
&=(1+\lambda+p)\,2^{-\mu-p}\,a^{\mu-1-\lambda-p}\,y^{p+1}\,\frac{\Gamma(2\mu)\,\Gamma(1+\lambda+p-\mu)}{\Gamma\left(\frac{3}{2}\right)\,\Gamma(2+\lambda+p+\mu)\,\Gamma\left(1+\frac{b}{2}+p\right)}\\
&\times {}_4F_5\left[\begin{matrix}\frac{3}{2}+\frac{\lambda}{2}+\frac{p}{2},\ \frac{1}{2}+\frac{\lambda}{2}+\frac{p}{2}-\frac{\mu}{2},\ 1+\frac{\lambda}{2}+\frac{p}{2}-\frac{\mu}{2},\ 1;\\ \frac{1}{2}+\frac{\lambda}{2}+\frac{p}{2},\ 1+\frac{\lambda}{2}+\frac{p}{2}+\frac{\mu}{2},\ \frac{3}{2}+\frac{\lambda}{2}+\frac{p}{2}+\frac{\mu}{2},\ \frac{1}{2}+p,\ \frac{3}{2};\end{matrix}\ -\frac{y^2}{4a^2}\right].
\end{aligned}\tag{3.4}$$

Corollary 3.4. *Let* $a,\ y \in \mathbb{R}^+$. *Also, let* $p,\ \lambda,\ \mu \in \mathbb{C}$ *with* $\Re(\mu+p) > -1$ *and* $\Re(\lambda) > \Re(\mu)$. *Then*

$$\begin{aligned}
&\int_0^\infty x^{\mu-1}\left(x+a+\sqrt{x^2+2ax}\right)^{-\lambda}\mathcal{W}_{p,b,c}\left(\frac{xy}{x+a+\sqrt{x^2+2ax}}\right)dx \\
&\quad = 2^{-\mu-2p-1}\,a^{\mu-\lambda}\,y^{p+1}\,\Gamma(\lambda-\mu) \\
&\times {}_3\Psi_4\left[\begin{matrix}(1,\,1)\,,\ (\lambda+p+2,\,2)\,,\ (2\mu+2p+2,\,4)\,; \\ \left(\frac{3}{2},\,1\right),\ \left(p+\frac{1}{2},\,1\right),\ (\lambda+p+1,\,2)\,,\ (\lambda+\mu+2p+3,\,4)\,;\end{matrix}\ -\frac{y^2}{16}\right].
\end{aligned} \tag{3.5}$$

References

[1] P. Agarwal , S. Jain, S. Agarwal and M. Nagpal, On a new class of integrals involving Bessel functions of the first kind, *Commun. Numer. Anal.* **2014** (2014), 7 pages.

[2] A. R. Ahmadi and S. E. Widnall, Unsteady lifting-line theory as a singular-perturbation problem, *J. Fluid Mech.* **153** (1985), 59–81.

[3] S. Ali, On some new unified integrals, *Adv. Comput. Math. Appl.* **1**(3) (2012), 151–153.

[4] J. Choi and P. Agarwal, Certain unified integrals associated with Bessel functions, *Bound. Value Probl.*, 2013 (2013), Artile ID 95, 9 pages.

[5] J. Choi, P. Agarwal, S. Mathur and S. D. Purohit, Certain new integral formulas involving the generalized Bessel functions, *Bull. Korean Math. Soc.* **51**(4) (2014), 995–1003.

[6] J. Choi, K. S. Nisar, S. Jain and P. Agarwal, Certain generalized Appell type functions and their properties, *Appl. Math. Sci.* **9**(132) (2015), 6567–6581.

[7] C. Fox, The asymptotic expansion of generalized hypergeometric functions, *Proc. London Math. Soc.* **27**(2) (1928), 389–400.

[8] A. Hasanov, H. M. Srivastava, Decomposition formulas associated with the Lauricella multivariable hypergeometric functions, *Comput. Math. Appl.* **53**(7) (2007), 1119–1128.

[9] M. H. Hirata, Flow near the bow of a steadily turning ship, *J. Fluid Mech.* **71**(2) (1975), 283–291.

[10] K. S. Nisar and S. R. Mondal, Certain unified integral formulas involving the generalized modified k-Bessel function of the first kind, arXiv:1601.06487 [math.CA].

[11] K. S. Nisar , S. D. Purohit and S. R. Mondal, Generalized fractional kinetic equations involving generalized Struve function of the first kind, *J. King Saud University-Sci.* **28**(2)(2016), 167-171.

[12] K. S. Nisar, R. K. Parmar and A. H. Abusufian, Certain new unified integrals associated with the generalized k-Bessel function, *Far East J. Math. Sci.* **100**(9) (2016), 1533–1544.

[13] F. Oberhettinger, *Tables of Mellin Transforms*, Springer-Verlag, New York, 1974.

[14] T. G. Pedersen, Variational approach to excitons in carbon nanotubes, *Phys. Rev. B* **67**(7) (2003), 073401.

[15] J. Shao and P. Hänggi, Decoherent dynamics of a two-level system coupled to a sea of spins, *Phys. Rev. Lett.* **81**(26) (1998), 5710–5713.

[16] D. C. Shaw, Perturbational results for diffraction of water-waves by nearly-vertical barriers, *IMA J. Appl. Math.* **34**(1) (1985), 99–117.

[17] D. P. Shukla, A note on generating relations for Lauricella's function of several variables, *Proc. Amer. Math. Soc.* **73**(2) (1979), 199–206.

[18] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.

[19] H. M. Srivastava and M. C. Daoust, A note on the convergence of Kampé de Fériet's double hypergeometric series, *Math. Nachr.* **53** (1985), 151–159.

[20] H. M. Srivastava, P. W. Karlsson, Multiple Gaussian Hypergeometric Series, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane, and Toronto, (1985).

[21] E. M. Wright, The asymptotic expansion of integral functions defined by Taylor series, *Philos. Trans. Roy. Soc. London, Ser. A.* **2**38 (1940), 423–451.

[22] E. M. Wright, The asymptotic expansion of the generalized hypergeometric function II, *Proc. London Math. Soc.* (2) **4**6 (1940), 389–408.

K. S. Nisar: Department of Mathematics, College of Arts and Science-Wadi Al dawaser, Prince Sattam bin Abdulaziz University, Riyadh region 11991, Saudi Arabia

E-mail address: ksnisar1@gmail.com, n.sooppy@psau.edu.sa